

Implementing Interventions to Improve Care for People with Obesity in 10 U.S. Health Care Organizations Elizabeth L. Ciemins, PhD, MPH, MA; Vaishali Joshi; Earlean Chambers, RN; Danielle F. Casanova, MBA; John K. Cuddeback, MD, PhD — AMGA, Alexandria, Virginia

Study Objective — Within a national, primary care-based, population health-focused, obesity learning collaborative, to identify the most impactful implemented strategies to improve care for people with obesity.

Background

- 71.6% U.S. adults with overweight or obesity; 39.8% have obesity.¹
- Annual medical costs for people with obesity are \$1,429 per person more than for people of normal weight.²
- A comprehensive program to treat obesity requires a combination of lifestyle (diet and exercise), medical, and surgical management. STOP Obesity Alliance's proposed Standard of Obesity Care³ should be considered.
- Reimbursement challenges persist in the treatment of obesity.
- Effective implementation strategies are needed, with attention to local context and both internal and external influencing factors.
- 68% of AMGA members surveyed in 2015 reported they had not implemented obesity guidelines. In response, AMGA conducted a learning collaborative to address obesity in primary care and studied the implementation process.

AMGA Obesity Care Model Collaborative

Objective: 3-year collaborative to define, pilot, and evaluate a framework and necessary components to address obesity in multi-specialty medical groups, academic medical centers, and integrated health systems



Methods

Design: Qualitative analysis of data collected during site visits

conducted as part of a national learning collaborative in which 10 U.S. health care organizations (HCOs) implemented programs to improve care for patients with obesity in primary care. A subset of Evidence for Researchers Implementing Change (ERIC) Strategies (46 of 73) were mapped to current interventions and rated by participating HCOs as having high, medium, or low impact.

Population: Providers and care teams caring for approximately 225,000 primary care patients with overweight or obesity seen at least annually at 10 HCOs from 10/1/2016 through 6/30/2019.

References

- 1. https://www.cdc.gov/obesity/data/adult.html; https://www.cdc.gov/nchs/fastats/obesity-overweight.htm
- 2. Finkelstein EA, Trogdon JG, Cohen JW, Dietz W. Health Affairs. 2009;28:w822–w831
- 3. Dietz WH, Gallagher C. *Obesity*. 23 June 2019.

Table 1. Strategies with Highest Perceived Impact*	
Necessary for Program Initiation	Community
Identify & prepare champions	
Identify early adopters	
Assess for readiness & identify barriers, facilitators	
Develop educational materials	Healthcare
Conduct educational meetings	Organization
Organize clinician implementation team meetings	+
Necessary for Program Success, Sustainability	
Develop & organize quality monitoring systems	
Recruit, designate, train for leadership	
Use train-the-trainer strategies; dynamic training	Care Team
Make training dynamic	
Conduct small tests of change, e.g., PDSA cycles	6767
Purposely reexamine implementation	
Stage implementation scale-up	Patient & Family
Intervene with patients to enhance uptake & adherence	
Audit & provide feedback (transparent reporting)	
Tailor strategies	UUUUU
* 100% ranked high by \geq 5 HCOs; 67% by \geq 6; 20% by \geq 8 HCOs	

100 /o latiked high by $\geq 5 \Pi COS$, 67 /o by ≥ 6 , 20 /o by $\geq 6 \Pi COS$

About AMGA

AMGA (American Medical Group Association) is a non-profit trade association representing 440 multispecialty medical groups and integrated delivery systems with a total of 175,000 full-time equivalent physicians. AMGA conducts national campaigns and disease-focused collaboratives, such as this one, focused on helping member organizations achieve the quadruple aim. AMGA collaboratives include benchmarking against other highperforming organizations to identify best practices and translate to other AMGA member organizations.

Implications for Policy or Practice

- A learning collaborative approach may be effective in designing
- based on the individual context of each organization.
- education, and EHR alerts all potentially contributed to improved outcomes.
- strategies while simultaneously impacting patient care. (See Figure 1.)
- medications. (See Figure 2.)
- the management of patients with overweight or obesity.

Results

Table 2. Highest Perceived Impact Interventions and Potential Measure(s) Impacted, One Health Care Organization Example



Weight Classes

Overweight:	$25 \le BMI < 30$	Obesity Class 2:	$35 \le BMI < 40$
Obesity Class 1:	$30 \le BMI < 35$	Obesity Class 3:	$BMI \ge 40$

comprehensive clinical programs for people with obesity, particularly when evidence-based non-pharmacologic interventions are lacking.

• Mapping interventions to a common set of implementation strategies may be useful in determining which interventions are most frequently associated with clinically successful programs, among geographically and structurally diverse HCOs.

• Strategies may then be shared with HCOs starting new primary care-based obesity programs, to guide the selection of successful interventions

• Shared learning between HCOs, as well as high-impact strategies such as provider and clinic champions, transparent reporting, provider and staff

• Focus on financial implications, e.g., HCC coding for BMI \geq 40 among MSSP or MA patients, may complement other obesity management

• Need dedicated resources to focus on insurance coverage, including attention to coding and negotiating with payers, particularly around

• A learning collaborative approach that both builds care models and monitors clinical and process outcomes over time is promising in addressing

Figure 1. Obesity Diagnoses Pre/Post Intervention, by Weight Class ($n \cong 50,000$)

	Class 1 O	besity	Class 2 C	Obesity	Class 3 Obesity	
100% –						86.9%
80% –				CO 10/		
60% –		42 00/		60.1%	75.2%	
40% –		42.8%	47.8%			
20% –	26.6%					
0% –	20.070					
	2018 Q1	2019 Q2	2018 Q1	2019 Q2	2018 Q1	2019 Q2

"Knowledge and best practices are meant to be shared, and these meetings allow us to do just that. This collaborative has exposed us to new ideas and strategies, many of which we may never have thought of on our own."

— Marianne Sumego, MD, Community Internal Medicine, Cleveland Clinic



Figure 2. Anti-obesity Prescriptions Pre/Post Intervention, Pilot vs. Entire **Organization, by Weight Class (n \cong 100,000)**